arXiv:gr-qc/0511125v1 23 Nov 2005

Matter Inheritance Symmetries of Spherically Symmetric Static Spacetimes

M. Sharif *
Department of Mathematics, University of the Punjab,
Quaid-e-Azam Campus Lahore-54590, PAKISTAN.

Abstract

In this paper we discuss matter inheritance collineations by giving a complete classification of spherically symmetric static spacetimes by their matter inheritance symmetries. It is shown that when the energy-momentum tensor is degenerate, most of the cases yield infinite dimensional matter inheriting symmetries. It is worth mentioning here that two cases provide finite dimensional matter inheriting vectors even for the degenerate case. The non-degenerate case provides finite dimensional matter inheriting symmetries. We obtain different constraints on the energy-momentum tensor in each case. It is interesting to note that if the inheriting factor vanishes, matter inheriting collineations reduce to be matter collineations already available in the literature. This idea of matter inheritance collineations turn out to be the same as homotheties and conformal Killing vectors are for the metric tensor.

Keywords : Matter Inheritance Symmetries, Spherically Symmetric Static Spacetimes.

---

*e-mail: msharif@math.pu.edu.pk

1 Introduction

Let (M, g) be a spacetime, where M is a smooth, connected, Hausdorff four-dimensional manifold and g is smooth Lorentzian metric of signature (+ - - -) defined on M. The manifold M and the metric g are assumed smooth (C^∞). We shall use the usual component notation in local charts, and a covariant derivative with respect to the symmetric connection Γ associated with the metric g will be denoted by a semicolon and a partial derivative by a comma. A smooth vector field ξ is said to generate a Matter inheritance collineation (MIC) [1-2] if it satisfies the following equation

$$\pounds_\xi T_{ab} = \psi(x^a) T_{ab}, \tag{1}$$

where $\pounds$ is the Lie derivative operator, ξ^a is the symmetry or collineation vector. If $\psi = 0$, then $\pounds_\xi T_{ab} = 0$ and ξ is said to preserve a matter symmetry [3] on M or is said to generate a matter collineation (MC). If ξ is a Killing vector (KV), then $\pounds_\xi T_{ab} = 0$ and hence any isometry is also a MC. The converse is not true, in general. The MIC Eq.(1) can be written in component form as

$$T_{ab,c}\xi^c + T_{ac}\xi^c_{,b} + T_{cb}\xi^c_{,a} = \psi T_{ab}, \quad (a, b, c = 0, 1, 2, 3). \tag{2}$$

A recent literature [3-13] shows a significant interest in the study of the various symmetries (in particular, Ricci and matter collineations). These symmetries arise in the exact solutions of Einstein's field equations

$$R_{ab} - \frac{1}{2} R g_{ab} \equiv G_{ab} = \kappa T_{ab}, \tag{3}$$

where κ is the gravitational constant, G_{ab} is the Einstein tensor, R_{ab} is the Ricci and T_{ab} is the matter (energy-momentum) tensor. Also, $R = g^{ab} R_{ab}$ is the Ricci scalar. We have assumed here that the cosmological constant $\Lambda = 0$.

There exists a large body of literature on classification of spacetimes according to their isometries and the groups admitted by them [14-17]. These investigations of symmetries played an important role in the classification of spacetimes, giving rise to many interesting results with useful applications. As curvature and Ricci tensors play a significant role in understanding the geometric structure of metrics, the energy-momentum tensor enables us to understand the physical structure of spacetimes.

Many of the difficulties encountered in studying such symmetries in general relativity are due to the fact that the associated vector fields may constitute infinite dimensional vector space. Problems of this nature are encountered very rear in the study of other symmetries. Unlike in the usual study of affine, conformal and projective symmetry where the initial assumption that ξ is C^3 (C^2 for affines) is sufficient to ensure that ξ is smooth (C^∞), the solution of (2) may be smooth, or C^m (but not C^{m+1}) for any M with $1 \leq m < \infty$. Hence the set of solutions of (2), whilst necessarily a vector space, is not necessarily a Lie algebra under the usual Lie bracket operation. Of course, if $\xi_{(1)}$ and $\xi_{(2)}$ satisfy (2) then so does $[\xi_{(1)}, \xi_{(2)}]$ provided it is C^1. The set of all smooth MCs on M is, of course, a Lie algebra.

In a recent paper [11], we have classified spherically symmetric spacetimes according to their MCs. A complete classification of static plane symmetric and cylindrically symmetric static spacetimes according to their MCs has been given in very recent papers [12] and [13] respectively. Some interesting consequences turn up. Moreover, Duggal [1-2] has proposed a relationship of the Ricci inheritace symmetry with conformal KV. This paper is devoted to classify spherically symmetric static spacetimes according to their MI symmetry. The rest of the paper is organized as follows. In the next section, we shall write down MIC equations. In section 3, we shall solve these equations for different possibilities to obtain MIC. Finally, we shall summarize the results in section 4.

2 Matter Inheritance Equations

The most general spherically symmetric static metric is given as

$$ds^2 = e^{\nu(r)}dt^2 - e^{\lambda(r)}dr^2 - e^{\mu(r)}d\Omega^2, \tag{4}$$

The non-vanishing components of the energy-momentum tensor T_{ab} are

$$\begin{aligned}
T_0 &= \frac{e^{\nu-\lambda}}{4}(2\mu'\lambda' - 3\mu'^2 - 4\mu'') + e^{\nu-\mu}, \\
T_1 &= \frac{\mu'^2}{4} + \frac{\mu'\nu'}{2} - e^{\lambda-\mu}, \\
T_2 &= \frac{e^{\mu-\lambda}}{4}(2\mu'' + \mu'^2 - \mu'\lambda' + \mu'\nu' + 2\nu'' + \nu'^2 - \nu'\lambda'), \\
T_3 &= T_2 \sin^2\theta.
\end{aligned} \tag{5}$$

We can write MIC Eqs.(2) in the expanded form as follows

$$T_{0,1}\xi^1 + 2T_0\xi^0_{,0} = \psi T_0, \tag{6}$$

$$T_0\xi^0_{,1} + T_1\xi^1_{,0} = 0, \tag{7}$$

$$T_0\xi^0_{,2} + T_2\xi^2_{,0} = 0, \tag{8}$$

$$T_0\xi^0_{,3} + \sin^2\theta T_2\xi^3_{,0} = 0, \tag{9}$$

$$T_{1,1}\xi^1 + 2T_1\xi^1_{,1} = \psi T_1, \tag{10}$$

$$T_1\xi^1_{,2} + T_2\xi^2_{,1} = 0, \tag{11}$$

$$T_1\xi^1_{,3} + \sin^2\theta T_2\xi^3_{,1} = 0, \tag{12}$$

$$T_{2,1}\xi^1 + 2T_2\xi^2_{,2} = \psi T_2, \tag{13}$$

$$T_2(\xi^2_{,3} + \sin^2\theta\xi^3_{,2}) = 0, \tag{14}$$

$$T_{2,1}\xi^1 + 2T_2\cot\theta\xi^2 + 2T_2\xi^3_{,3} = \psi T_2. \tag{15}$$

For the sake of simplicity, we are using the notation $T_{aa} = T_a$.

3 Matter Inheritance Symmetries

In this section, we solve the MIC equations given by Eq.(6)-(15) for different situations to find MIC.

From Eq.(14), we have the following two possibilities:

$$\begin{array}{ll}(1) & T_2 = 0, \quad \xi^2_{,3} + \xi^3_{,2}\sin^2\theta \neq 0,\\ (2) & T_2 \neq 0, \quad \xi^2_{,3} + \xi^3_{,2}\sin^2\theta = 0.\end{array}$$

It is to be noted that the case (1) corresponds to degenerate case while in the case (2), some of the subcases will lead to non-degenerate case.

Case (1): When we use constraint of this case in MIC equations, we get

$$T_0\xi^0_{,2} = 0, \quad T_0\xi^0_{,3} = 0, \quad T_1\xi^1_{,2} = 0, \quad T_1\xi^1_{,3} = 0. \tag{16}$$

The last two equations give rise to the following two subcases:

$$\begin{array}{ll}(a_1) & T_1 = 0, \quad \xi^1_{,2} \neq 0 \neq \xi^1_{,3},\\ (a_2) & T_1 \neq 0, \quad \xi^1_{,2} = 0 = \xi^1_{,3}.\end{array}$$

Subcase 1(a_1): The subcase $1(a_1)$ yields further two possibilities:

$$\begin{array}{ll}(b_1) & T_0 = 0, \quad \xi^0_{,2} \neq 0 \neq \xi^0_{,3} \neq \xi^0_{,1}, \\ (b_2) & T_0 \neq 0, \quad \xi^0_{,2} = 0 = \xi^0_{,3} = \xi^0_{,1}.\end{array}$$

The first possibility $1a_1(b_1)$ gives arbitrary MI vectors while the second possibility $1a_1(b_2)$ leads to a contradiction.

Subcase 1(a_2): When we solve MIC equations for the subcase $1(a_2)$, we obtain the following two options

$$\begin{array}{ll}(b_1) & T_0 = 0, \quad \xi^0_{,2} \neq 0 \neq \xi^0_{,3}, \\ (b_2) & T_0 \neq 0, \quad \xi^0_{,2} = 0 = \xi^0_{,3}.\end{array}$$

In the first case $1a_2(b_1)$, i.e., $T_0 = 0,\ \xi^0_{,2} \neq 0 \neq \xi^0_{,3}$, MIC equations yield the following solution

$$\begin{aligned} \xi^l &= \xi^l(x^a), \quad l = 0, 2, 3, \\ \xi^1 &= \frac{1}{\sqrt{T_1}}\left(\frac{\psi}{2}\int \sqrt{T_1}dr + k\right), \end{aligned} \tag{17}$$

where k is an integration constant.

For the option $1a_2(b_2)$, simultaneous solution of MIC equations gives the following constraint

$$\frac{B_{,000}}{\dot{B}} = k = \frac{T_0}{\sqrt{T_1}}\left(\frac{T_0'}{2T_0\sqrt{T_1}}\right)' \tag{18}$$

where B is an arbitrary integration function of t and k is a separation constant. $B_{,0}$ and $\dot{B}$ are the same notations and represent derivative with respect to time. This gives us three more possibilities according to

$$(c_1) \quad k > 0, \quad (c_2) \quad k = 0, \quad (c_3) \quad k < 0.$$

In the first case $1a_2b_2(c_1)$, we obtain the following solution

$$\begin{aligned} \xi^0 &= \frac{\psi}{2}t - \frac{\psi T_0'}{4T_0\sqrt{T_1}}\int \sqrt{T_1}dr \\ &- \frac{T_0'}{2T_0k\sqrt{T_1}}[kc_0t + c_1\cosh\sqrt{k}t - c_2\sinh\sqrt{k}t] + c_3, \end{aligned}$$

$$\begin{aligned}
\xi^1 &= \frac{\psi}{2\sqrt{T_1}}\int\sqrt{T_1}dr \\
&+ \frac{1}{\sqrt{T_1}}[c_0+c_1\cosh\sqrt{k}t+c_2\sinh\sqrt{k}t]+c_3, \\
\xi^2 &= \xi^2(x^a), \quad \xi^3=\xi(x^a),
\end{aligned} \tag{19}$$

where

$$(\frac{T_0'}{4T_0\sqrt{T_1}})'=-k\frac{\sqrt{T_1}}{T_0}, \quad (\frac{\psi T_0'}{4T_0\sqrt{T_1}}\int\sqrt{T_1}dr)'=-c_0k\frac{\sqrt{T_1}}{T_0}.$$

For the case $1a_2b_2(c_2)$, Eq.(18) yields

$$B(t)=k_1\frac{t^2}{2}+k_2t+k_3, \quad \frac{T_0'}{2T_0\sqrt{T_1}}=m, \tag{20}$$

where m is an arbitrary constant which gives further three options according to

$$(d_1) \quad m>0, \quad (d_2) \quad m=0, \quad (d_3) \quad m<0.$$

The possibilities $1a_2b_2c_2(d_1)$ and $1a_2b_2c_2(d_3)$ lead to contradiction. The possibility $1a_2b_2c_2(d_2)$ gives the following MI vectors

$$\begin{aligned}
\xi^0 &= \frac{\psi}{2}t+k_4, \\
\xi^1 &= \frac{\psi}{2}t-\int\sqrt{T_1}dr+\frac{k_3}{\sqrt{T_1}}, \\
\xi^j &= \xi^j(x^a), \quad j=2,3.
\end{aligned} \tag{21}$$

The case $1a_2b_2(c_3)$ gives the similar results as the case $1a_2b_1(c_1)$. We see that all the possibilities of the case 1 yield infinite dimensional MIC as it is expected for the degenerate energy-momentum tensor.

Case 2: Solving MIC Eqs.(11), (12), (16) and (17) simultaneously, we obtain

$$T_1(\xi^1_{,23}-\cot\theta\xi^1_{,3})=0, \tag{22}$$

From here, it follows that either

$$\begin{aligned}
&(a_1) \quad T_1=0, \quad \xi^1_{,23}-\xi^1_{,3}\cot\theta\neq 0, \\
or \quad &(a_2) \quad T_1\neq 0, \quad \xi^1_{,23}-\xi^1_{,3}\cot\theta=0.
\end{aligned}$$

Subcase 2(a_1): In this case, it is obvious from MIC Eq.(7) that we have the following two possibilities:

$$\begin{array}{ll}(b_1) & T_0=0, \quad \xi^0_{,1}\neq 0,\\ (b_2) & T_0\neq 0, \quad \xi^0_{,1}=0.\end{array}$$

When we consider the case $2a_1(b_1)$, we obtain the following solution

$$\begin{aligned}\xi^0 &= \xi^0(x^a),\\ \xi^i &= \xi^i(\theta,\phi), \quad i=1,2,3.\end{aligned} \tag{23}$$

For the case $2a_1(b_2)$, when we solve MIC equations, we arrive at the following result

$$(\frac{T_0}{T_2})'\xi^0_{,2}=0$$

which yields the following two options

$$\begin{array}{ll}(c_1) & (\frac{T_0}{T_2})'=0, \quad \xi^0_{,2}\neq 0,\\ (c_2) & (\frac{T_0}{T_2})'\neq 0, \quad \xi^0_{,2}=0.\end{array}$$

In the case $2a_1b_2(c_1)$, solution of MIC equations gives the following MI vectors

$$\begin{aligned}\xi_{(0)} &= \partial_t,\\ \xi_{(1)} &= \sin\phi\partial_\theta+\cot\theta\cos\phi\partial_\phi,\\ \xi_{(2)} &= \cos\phi\partial_\theta-\cot\theta\sin\phi\partial_\phi,\\ \xi_{(3)} &= \partial_\phi,\\ \xi_{(4)} &= \psi\frac{T_0}{T_0'}\partial_r,\\ \xi_{(5)} &= k_1(\cos t\partial_t+\frac{2T_0}{T_0'}\sin t\partial_r)\sin\theta\sin\phi\\ &- (\cos\theta\sin\phi\partial_\theta+\csc\theta\cos\phi\partial_\phi)\cos t,\\ \xi_{(6)} &= k_1(\sin t\partial_t-\frac{2T_0}{T_0'}\cos t\partial_r)\sin\theta\sin\phi\\ &- (\cos\theta\sin\phi\partial_\theta+\csc\theta\cos\phi\partial_\phi)\sin t,\end{aligned}$$

$$\begin{aligned}
\xi_{(7)} &= k_1(\cos t\partial_t + \frac{2T_0}{T_0'}\sin t\partial_r)\sin\theta\cos\phi \\
&- (\cos\theta\cos\phi\partial_\theta + \csc\theta\sin\phi\partial_\phi)\cos t, \\
\xi_{(8)} &= k_1(\sin t\partial_t - \frac{2T_0}{T_0'}\cos t\partial_r)\sin\theta\cos\phi \\
&- (\cos\theta\cos\phi\partial_\theta + \csc\theta\sin\phi\partial_\phi)\sin t, \\
\xi_{(9)} &= k_1(\cos t\partial_t + \frac{2T_0}{T_0'}\sin t\partial_r)\cos\theta + \cos t\sin\theta\partial_\theta, \\
\xi_{(10)} &= k_1(\sin t\partial_t - \frac{2T_0}{T_0'}\cos t\partial_r)\cos\theta + \sin t\sin\theta\partial_\theta. \qquad (24)
\end{aligned}$$

Thus we obtain eleven independent MI vectors in this degenerate case in which four are the usual isometries of the spherically symmetric spacetime.

The case $2a_1b_2(c_2)$ yields the following five independent MIC

$$\begin{aligned}
\xi_{(0)} &= (\frac{\psi(k_2-1)}{2k_2}t+1)\partial_t, \\
\xi_{(1)} &= \sin\phi\partial_\theta + \cot\theta\cos\phi\partial_\phi, \\
\xi_{(2)} &= \cos\phi\partial_\theta - \cot\theta\sin\phi\partial_\phi, \\
\xi_{(3)} &= \partial_\phi, \\
\xi_{(4)} &= \psi\frac{T_0}{T_0'}\partial_r. \qquad (25)
\end{aligned}$$

where $T_2' \neq 0,\ T_2 = k_1 T_0^{k_2},\ k_1 \neq 0,\ k_2 \neq 0, 1.$ Thus we obtain finite dimensional MICs in another degenerate case.

Subcase 2(a_2)**:** When we solve MIC Eqs.(8), (9), (14) and (20) together with the constraint of this case, we have

$$T_0(\xi^0_{,23} - \cot\theta\xi^0_{,3}) = 0, \qquad (26)$$

From this equation, we further have the following two possibilities:

$$\begin{aligned}
(b_1) \quad T_0 = 0, \quad \xi^0_{,23} - \xi^0_{,3}\cot\theta \neq 0, \\
(b_2) \quad T_0 \neq 0, \quad \xi^0_{,23} - \xi^0_{,3}\cot\theta = 0.
\end{aligned}$$

For the case $2a_2(b_1)$, when we solve MIC equations simultaneously for these constraints, we obtain the following solution

$$\xi_{(0)} = \xi^0(x^a)\partial_t,$$

$$\begin{aligned}
\xi_{(1)} &= \sin\phi\partial_\theta + \cot\theta\cos\phi\partial_\phi, \\
\xi_{(2)} &= \cos\phi\partial_\theta - \cot\theta\sin\phi\partial_\phi, \\
\xi_{(3)} &= \partial_\phi, \\
\xi_{(4)} &= \psi\frac{T_2}{T_2'}, \\
\xi_{(5)} &= \frac{2\sqrt{T_2}}{T_2'}\sin\theta\sin\phi\partial_r + \frac{1}{\sqrt{T_2}}\cos\theta\sin\phi\partial_\theta + \frac{1}{\sqrt{T_2}}\csc\theta\cos\phi\partial_\phi, \\
\xi_{(6)} &= \frac{2\sqrt{T_2}}{T_2'}\sin\theta\cos\phi\partial_r + \frac{1}{\sqrt{T_2}}\cos\theta\cos\phi\partial_\theta + \frac{1}{\sqrt{T_2}}\csc\theta\sin\phi\partial_\phi, \\
\xi_{(7)} &= \frac{2\sqrt{T_2}}{T_2'}\cos\theta\partial_r - \frac{1}{\sqrt{T_2}}\sin\theta\partial_\theta, \qquad (27)
\end{aligned}$$

where $4T_1T_2 = T_2'^2$. This gives infinite dimensional MICs.

The case $2a_2(b_2)$, i.e., $T_0 \neq 0, \quad \xi^0_{,23} - \xi^0_{,3}\cot\theta = 0$ turns out to be non-degenerate. When we use these constraints in the MIC Eqs.(6)-(15) and solve them simultaneously, after some tedious algebra, we arrive at the following solution

$$\xi^0 = -\frac{T_2}{T_0}[(\dot{A}_1\sin\phi - \dot{A}_2\cos\phi)\sin\theta + \dot{A}_3\cos\theta] + A_4, \qquad (28)$$

$$\xi^1 = -\frac{T_2}{T_1}[(A_1'\sin\phi - A_2'\cos\phi)\sin\theta + A_3'\cos\theta] + A_5, \qquad (29)$$

$$\xi^2 = (A_1\sin\phi - A_2\cos\phi)\cos\theta + A_3\sin\theta + c_1\sin\phi - c_2\cos\phi, \qquad (30)$$

$$\xi^3 = (A_1\cos\phi + A_2\sin\phi)\mathrm{cosec}\theta + (c_1\cos\phi + c_2\sin\phi)\cot\theta + c_0, \qquad (31)$$

where c_0, c_1 and c_2 are arbitrary constants and $A_\mu = A_\mu(t,r),\ \mu = 1,2,3,4,5$ are arbitrary functions. Here dot and prime indicate the differentiation with respect to time and r coordinate respectively. When we substitute these values of ξ^a in MIC Eqs.(6)-(15), we obtain the following differential constraints on A_ν with $c_0 = 0$. These ξ^a are satisfied subject to the following differential constraints on A_μ

$$2T_1\ddot{A}_i + T_{0,1}A_i' = 0, \quad i = 1,2,3, \qquad (32)$$

$$2T_0\dot{A}_4 + T_{0,1}A_5 = \psi T_0, \qquad (33)$$

$$2T_2\dot{A}_i' + T_0\left(\frac{T_2}{T_0}\right)'\dot{A}_i = 0, \qquad (34)$$

$$T_0A'_4+T_1\dot{A}_5=0, \tag{35}$$

$$\left\{T_{1,1}\frac{T_2}{T_1}+2T_1\left(\frac{T_2}{T_1}\right)'\right\}A'_i+2T_2A''_i=0, \tag{36}$$

$$T_{1,1}A_5+2T_1A'_5=\psi T_1, \tag{37}$$

$$T_{2,1}A'_i+2T_1A_i=0,\quad c_0=0, \tag{38}$$

$$T_{2,1}A_5=\psi T_2. \tag{39}$$

Now the problem of working out MICs for all possibilities of A_i, A_4, A_5 is reduced to solving the set of Eqs.(28)-(31) subject to the above constraints. We start the classification of MICs by considering the constraint Eq.(39). It follows that A_5 is a function of coordinate r only and

$$\psi=\frac{T'_2A_5(r)}{T_2}. \tag{40}$$

When we use this value in Eqs.(33), (35) and (37) and solve simultaneously, we obtain

$$A_4(t)=\frac{k}{2}[\frac{T_0T'_2-T'_0T_2}{T_0\sqrt{T_1T_2}}]t+c_0, \tag{41}$$

where k is an integration constant. When we differentiate this equation with respect to time, we have

$$T_0T'_2-T'_0T_2=mT_0\sqrt{T_1T_2}, \tag{42}$$

where m is an arbitrary constant which can be zero or non-zero.

$$(c_1)\quad m=0,\quad (c_2)\quad m\neq 0.$$

The case $2a_2b_2(c_1)$ gives

$$A_i=(\frac{a_it^2}{2}+b_it)-a_i\int\frac{2T_1}{T'_0}dr+c_i. \tag{43}$$

Now if we use Eqs.(38), (40) and (41), we finally obtain the following MIC

$$\begin{aligned}
\xi_{(0)} &= \partial_t,\\
\xi_{(1)} &= \sin\phi\partial_\theta+\cot\theta\cos\phi\partial_\phi,\\
\xi_{(2)} &= \cos\phi\partial_\theta-\cot\theta\sin\phi\partial_\phi,\\
\xi_{(3)} &= \partial_\phi,\\
\xi_{(4)} &= \psi\frac{T_0}{T'_0} \qquad (44)
\end{aligned}$$

which gives five MI vectors.

For the case $2a_2b_2(c_2)$ when $m \neq 0$, we use Eqs.(32), (41) and (42) to obtain

$$\frac{\ddot{B}_i(t)}{B_i(t)} = -\frac{T_0'(\sqrt{\frac{T_0}{T_2}})'}{2T_2\sqrt{\frac{T_0}{T_2}}} = \alpha_i, \qquad (45)$$

where α_i is a separation constant which yields the following three different possibilities:

$$(d_1) \quad \alpha_i > 0, \quad (d_2) \quad \alpha_i = 0, \quad (d_3) \quad \alpha_i < 0.$$

In the case $2a_2b_2c_2(d_1)$, i.e., $\alpha_i > 0$, when we solve Eqs.(32), (34), (36) and (38), we obtain

$$[\frac{T_1'}{2T_1} + \frac{T_1}{T_2}(\frac{T_2}{T_1})'](\sqrt{\frac{T_0}{T_2}})' + (\sqrt{\frac{T_0}{T_2}})''[(a_i \sinh\sqrt{\alpha_i}t + b_i \cosh\sqrt{\alpha_i}t\frac{1}{\sqrt{\alpha_i}}] = 0 \quad (46)$$

It follows from the above equation that either

$$\begin{array}{ll} (e_1) & [\frac{T_1'}{2T_1} + \frac{T_1}{T_2}(\frac{T_2}{T_1})'](\sqrt{\frac{T_0}{T_2}})' + (\sqrt{\frac{T_0}{T_2}})'' = 0; \quad a_i \neq 0 \neq b_i, \\ or \quad (e_2) & [\frac{T_1'}{2T_1} + \frac{T_1}{T_2}(\frac{T_2}{T_1})'](\sqrt{\frac{T_0}{T_2}})' + (\sqrt{\frac{T_0}{T_2}})'' \neq 0; \quad a_i = 0 = b_i. \end{array}$$

However, both the above cases $2a_2b_2c_2d_1(e_1)$ and $2a_2b_2c_2d_1(e_2)$ give contradiction and hence can be excluded.

In the case $2a_2b_2c_2(d_2)$, i.e., $\alpha_i = 0$ when we use constraint equations, we obtain the following MIC

$$\begin{aligned}
\xi_{(0)} &= \partial_t, \\
\xi_{(1)} &= \sin\phi\partial_\theta + \cot\theta\cos\phi\partial_\phi, \\
\xi_{(2)} &= \cos\phi\partial_\theta - \cot\theta\sin\phi\partial_\phi, \\
\xi_{(3)} &= \partial_\phi, \\
\xi_{(4)} &= \sqrt{T_2}\sin\theta\sin\phi\partial_t - \frac{2\sqrt{T_2}}{T_2'}t\sin\theta\sin\phi\partial_r - \frac{1}{\sqrt{T_2}}t\cos\theta\sin\phi\partial_\theta, \\
\xi_{(5)} &= \sqrt{T_2}\sin\theta\cos\phi\partial_t - \frac{2\sqrt{T_2}}{T_2'}t\sin\theta\cos\phi\partial_r - \frac{1}{\sqrt{T_2}}t\cos\theta\cos\phi\partial_\theta,
\end{aligned}$$

$$
\begin{aligned}
\xi_{(6)} &= \sqrt{T_2}\cos\theta\partial_t - \frac{2\sqrt{T_2}}{T_2'}t\cos\theta\partial_r + \frac{1}{\sqrt{T_2}}t\sin\theta\partial_\theta, + \frac{1}{\sqrt{T_2}}t\csc\theta\cos\phi\partial_\phi, \\
\xi_{(7)} &= \frac{2\sqrt{T_2}}{T_2'}\sin\theta\sin\phi\partial_r + \frac{1}{\sqrt{T_2}}\cos\theta\sin\phi\partial_\theta + \frac{1}{\sqrt{T_2}}\csc\theta\cos\phi\partial_\phi, \\
\xi_{(8)} &= \frac{2\sqrt{T_2}}{T_2'}\sin\theta\cos\phi\partial_r + \frac{1}{\sqrt{T_2}}\cos\theta\cos\phi\partial_\theta + \frac{1}{\sqrt{T_2}}\csc\theta\sin\phi\partial_\phi, \\
\xi_{(9)} &= \frac{2\sqrt{T_2}}{T_2'}\cos\theta\partial_r - \frac{1}{\sqrt{T_2}}\sin\theta\partial_\theta, \\
\xi_{(10)} &= \psi\frac{T_2}{T_2'}\partial_r \qquad (47)
\end{aligned}
$$

which gives eleven MI vectors.

The case $2a_2b_2c_2(d_3)$, i.e., $\alpha_i < 0$ gives similar results as the case $2a_2b_2c_2(d_1)$.

4 Conclusion

We know that when the spherically symmetric static spacetimes are classified according to their MCs [11], we obtain either *four*, *six*, *seven* or *ten* independent MCs for the non-degenerate energy-momentum tensor. For the degenerate case, we obtain infinite dimensional MCs except the two cases where the energy-momentum tensor is degenerate but the group of matter collineations is finite-dimensional, i.e. *four* or *ten*.

In this paper, we have classified spherically symmetric spcetimes according to their MICs. We see that for the degenerate energy-momentum tensor, most of the cases yield infinite dimensional MICs. The worth mentioning cases are those where we have got finite number of MICs even when the energy-momentum tensor is zero. We obtain two such different cases having either eleven or five independent MICs. In the non-degenerate case, we obtain either five or eleven independent MICs. These contain the usual four isometries of the spherically symmetry and the rest are the proper (non-trivial) MICs. It can easily be seen from these cases that if we take the inheriting factor $\psi = 0$, MICs reduce to MCs. These MCs exactly coincide with those already found in the literature [11]. It is mentioned here that MICs coincide with RICs [18] but the constraint equations are completely different. We conclude that the idea of matter inheritance collineations turns out to be the same as homotheties and conformal Killing vectors for the metric tensor. The results are summarized in the form of tables given below.

Table 1. MICs for Spherically Symmetric Static Spacetimes

Cases	**MICs**	**Constraints**
$1a_1(b_1)$	Infinte No. of MICs	$T_2=0,\ \xi^2_{,3}+\xi^3_{,2}\sin^2\theta\neq 0,\ T_1=0,$ $\xi^1_{,2}\neq 0\neq \xi^1_{,3},\ T_0=0,$ $\xi^0_{,2}\neq 0\neq \xi^0_{,3}\neq \xi^0_{,1}$
$1a_1(b_2)$	Contradiction	$T_2=0,\ \xi^2_{,3}+\xi^3_{,2}\sin^2\theta\neq 0,\ T_1=0,$ $\xi^1_{,2}\neq 0\neq \xi^1_{,3},\ T_0\neq 0,$ $\xi^0_{,2}=0=\xi^0_{,3}=\xi^0_{,1}$
$1a_2(b_1)$	Infinte No. of MICs	$T_2=0,\ \xi^2_{,3}+\xi^3_{,2}\sin^2\theta\neq 0,\ T_1\neq 0,$ $\xi^1_{,2}=0=\xi^1_{,3},\ T_0=0,\ \xi^0_{,2}\neq 0\neq \xi^0_{,3}$
$1a_2b_2(c_1)$	Infinte No. of MICs	$T_2=0,\ \xi^2_{,3}+\xi^3_{,2}\sin^2\theta\neq 0,\ T_1\neq 0,$ $\xi^1_{,2}=0=\xi^1_{,3},\ T_0\neq 0,\ \xi^0_{,2}=0=\xi^0_{,3},$ $\frac{T_0}{T_1}(\frac{T_0'}{2T_0\sqrt{T_1}})'=k,\ k>0$
$1a_2b_2c_2(d_1)$	Contradiction	$T_2=0,\ \xi^2_{,3}+\xi^3_{,2}\sin^2\theta\neq 0,\ T_1\neq 0,$ $\xi^1_{,2}=0=\xi^1_{,3},\ T_0\neq 0,\ \xi^0_{,2}=0=\xi^0_{,3},$ $k=0,\frac{T_0'}{2T_0\sqrt{T_1}}=m,\ m>0$
$1a_2b_2c_2(d_2)$	Infinite No. of MICs	$T_2=0,\ \xi^2_{,3}+\xi^3_{,2}\sin^2\theta\neq 0,\ T_1\neq 0,$ $\xi^1_{,2}=0=\xi^1_{,3},\ T_0\neq 0,\ \xi^0_{,2}=0=\xi^0_{,3},$ $k=0,\ m=0$
$1a_2b_2c_2(d_3)$	Contradiction	$T_2=0,\ \xi^2_{,3}+\xi^3_{,2}\sin^2\theta\neq 0,\ T_1\neq 0,$ $\xi^1_{,2}=0=\xi^1_{,3},\ T_0\neq 0,\ \xi^0_{,2}=0=\xi^0_{,3},$ $k=0,\ m<0$
$2a_1(b_1)$	Infinite No. of MICs	$T_2\neq 0,\ \xi^2_{,3}+\xi^3_{,2}\sin^2\theta=0,\ T_1=0,$ $\xi^1_{,23}-\xi^1_{,3}\cot\theta\neq 0,\ T_0=0,\ \xi^0_{,1}\neq 0,$
$2a_1b_2(c_1)$	11	$T_2\neq 0,\ \xi^2_{,3}+\xi^3_{,2}\sin^2\theta=0,\ T_1=0,$ $\xi^1_{,23}-\xi^1_{,3}\cot\theta\neq 0,\ T_0\neq 0,\ \xi^0_{,1}=0,$ $(\frac{T_0}{T_2})'=0,\ \xi^0_{,2}\neq 0$
$2a_1b_2(c_2)$	5	$T_2\neq 0,\ \xi^2_{,3}+\xi^3_{,2}\sin^2\theta=0,\ T_1=0,$ $\xi^1_{,23}-\xi^1_{,3}\cot\theta\neq 0,\ T_0\neq 0,\ \xi^0_{,1}=0,$ $(\frac{T_0}{T_2})'\neq 0,\ \xi^0_{,2}=0$
$2a_2(b_1)$	Infinte No. of MICs	$T_2\neq 0,\ \xi^2_{,3}+\xi^3_{,2}\sin^2\theta=0,$ $T_1\neq 0,\ \xi^1_{,23}-\xi^1_{,3}\cot\theta=0,$ $T_0=0,\ \xi^0_{,23}-\xi^0_{,3}\cot\theta\neq 0$

Cases	**MICs**	**Constraints**
$2a_2b_2(c_1)$	5	$T_2 \neq 0,\ \xi^2_{,3} + \xi^3_{,2}\sin^2\theta = 0,$ $T_1 \neq 0,\ \xi^1_{,23} - \xi^1_{,3}\cot\theta = 0,$ $T_0 \neq 0,\ \xi^0_{,23} - \xi^0_{,3}\cot\theta = 0,$ $T_0T_2' - T_0'T_2 = mT_0\sqrt{T_1T_2},\ m = 0$
$2a_2b_2c_2d_1(e_1)$	Contradiction	$T_2 \neq 0,\ \xi^2_{,3} + \xi^3_{,2}\sin^2\theta = 0,$ $T_1 \neq 0,\ \xi^1_{,23} - \xi^1_{,3}\cot\theta = 0,$ $T_0 \neq 0,\ \xi^0_{,23} - \xi^0_{,3}\cot\theta = 0,$ $m \neq 0,\ \ -\frac{T_0'(\sqrt{\frac{T_0}{T_2}})'}{2T_2\sqrt{T_0}T_2} = \alpha_i,\ \alpha_i > 0,$ $(\frac{T_1'}{2T_1} + \frac{T_1}{T_2}(\frac{T_2}{T_1})')(\sqrt{\frac{T_0}{T_2}})'$ $+(\sqrt{\frac{T_0}{T_2}})'' = 0,\ a_i \neq 0 \neq b_i$
$2a_2b_2c_2d_1(e_2)$	Contradiction	$T_2 \neq 0,\ \xi^2_{,3} + \xi^3_{,2}\sin^2\theta = 0,$ $T_1 \neq 0,\ \xi^1_{,23} - \xi^1_{,3}\cot\theta = 0,$ $T_0 \neq 0,\ \xi^0_{,23} - \xi^0_{,3}\cot\theta = 0,$ $m \neq 0,\ \alpha_i > 0,$ $(\frac{T_1'}{2T_1} + \frac{T_1}{T_2}(\frac{T_2}{T_1})')(\sqrt{\frac{T_0}{T_2}})'$ $+(\sqrt{\frac{T_0}{T_2}})'' \neq 0,\ a_i = 0 = b_i$
$2a_2b_2c_2(d_2)$	11	$T_2 \neq 0,\ \xi^2_{,3} + \xi^3_{,2}\sin^2\theta = 0,$ $T_1 \neq 0,\ \xi^1_{,23} - \xi^1_{,3}\cot\theta = 0,$ $T_0 \neq 0,\ \xi^0_{,23} - \xi^0_{,3}\cot\theta = 0,$ $m \neq 0,\ \alpha_i = 0$
$2a_2b_2c_2(d_3)$	$2a_2b_2c_2(d_1)$	$T_2 \neq 0,\ \xi^2_{,3} + \xi^3_{,2}\sin^2\theta = 0,$ $T_1 \neq 0,\ \xi^1_{,23} - \xi^1_{,3}\cot\theta = 0,$ $T_0 \neq 0,\ \xi^0_{,23} - \xi^0_{,3}\cot\theta = 0,$ $m \neq 0,\ \alpha_i < 0$

It follows from these tables that each case has different constraints on the energy-momentum tensor. It would be interesting to solve these constraints to ascertain the non-trivial MIC.

References

[1] Duggal, K.L.: J. Math. Phys., **33**(1992)2989.

[2] Duggal, K.L.: Acta Applicanade Mathematica, **31**(1993)225.

[3] Hall, G.S., Roy, I. and Vaz, L.R.: Gen. Rel and Grav. **28**(1996)299.

[4] Camcı, U. and Barnes, A.: Class. Quant. Grav. **19**(2002)393.

[5] Carot, J. and da Costa, J.: *Procs. of the 6th Canadian Conf. on General Relativity and Relativistic Astrophysics*, Fields Inst. Commun. 15, Amer. Math. Soc. WC Providence, RI(1997)179.

[6] Carot, J., da Costa, J. and Vaz, E.G.L.R.: J. Math. Phys. **35**(1994)4832.

[7] Sharif, M.: Nuovo Cimento **B116**(2001)673;
Astrophys. Space Sci. **278**(2001)447.

[8] Camcı, U. and Sharif, M.: Gen Rel. and Grav. **35**(2003)97.

[9] Camcı, U. and Sharif, M.: Class. Quant. Grav. **20**(2003)2169-2179.

[10] Tsamparlis, M. and Apostolopoulos, P.S.: Gen. Rel. and Grav. **36**(2004)47.

[11] Sharif, M. and Sehar Aziz: Gen Rel. and Grav. **35**(2003)1091;
Sharif, M.: J. Math. Phys. **44**(2003)5142.

[12] Sharif, M.: J. Math. Phys. **45**(2004)1518;

[13] Sharif, M.: J. Math. Phys. 45(2004)1532.

[14] Katzin, G.H., Levine J. and Davis, W.R.: J. Math. Phys. **10**(1969)617.

[15] Petrov, A.Z.: *Einstein Spaces* (Pergamon, Oxford University Press, 1969).

[16] Stephani, H., Kramer, D., MacCallum, M.A.H., Hoenselaers, C. and Hearlt, E.: *Exact Solutions of Einstein's Field Equations* (Cambridge University Press, 2003).

[17] Rcheulishrili, G.: J. Math. Phys. **33**(1992)1103.

[18] Bokhari, A.H., Kashif, A.R. and Kara A.H.: Nuovo Cimento **118B**(2003)803.